arXiv:nlin/0602015v1 [nlin.SI] 8 Feb 2006

Constructing periodic wave solutions of nonlinear equations by Hirota bilinear method

Huihui Dai[a], Engui Fan[b], Xianguo Geng[c]

a. Department of Mathematics, City University of Hong Kong, Hong Kong SAR, China

b. Institute of Mathematics, Fudan University, Shanghai 200433, P. R. China.

c. Department of Mathematics, Zhengzhou University, Zhengzhou, P. R. China

Abstract. Hirota bilinear method is proposed to directly construct periodic wave solutions in terms of Riemann theta functions for nonlinear equations. The asymptotic property of periodic waves are analyzed in detail. It is shown that well-known soliton solutions can be reduced from the periodic wave solutions.

1. Introduction

The investigation of the exact solutions of nonlinear equations plays an important role in the study of nonlinear physical phenomena. For example, the wave phenomena observed in fluid dynamics, plasma and elastic media are often modelled by the bell shaped sech solutions and the kink shaped tanh traveling wave solutions. The exact solution, if available, of those nonlinear equations facilitates the verification of numerical solvers and aids in the stability analysis of solutions. In the past decades, there has been significant progression in the development of these methods such as inverse scattering method [1,2], Darboux transformation [3-6], Hirota bilinear method [7-12], algebro-geometric method [13-18] and others. Among them, the algebro-geometric method is an analogue of inverse scattering transformation, which was first developed by Matveev, Its, Novikov, and Dubrovin et al. The method can derive an important class of exact solutions, which is called quasi-periodic or algebro-geometric solution, to many soliton equations such as KdV equation, sin-Gordon equation, and Schrodinger equation. In recent years, such solutions of nonlinear equations have been aroused much interest in the mathematical physics [19-24]. However, this method usually is applied in the integrable nonlinear evolution equations admitting Lax pairs representation and involves complicated algebraic geometry theory. These have been used far less than their soliton counterparts. The main reason for this is that they are far more complicated. Soliton solutions are typically expressed in terms of rational or hyperbolic functions, whereas qusi-periodic solutions require the use of Riemann theta functions and calculus on Riemann surfaces. Recently, Deconinck, Hoeij et al proposed an algorithm to compute the Riemann theta function and Riemann constants [25,26].

It is well known that the bilinear derivative method developed by Hirota is a powerful and direct approach to construct exact solution of nonlinear equations [7-9]. Once a nonlinear equation is written in bilinear forms by a dependent variable transformation, then multi-soliton solutions and rational solutions can be obtained. In recent years, Hirota method also has been developed for obtaining Wronskian and Pfaffian forms of N-soliton solution [10,11]. In this paper, we propose Hirota method for directly constructing periodic wave solutions in Riemann theta functions. It is shown that the periodic solutions can be reduced to classical soliton solutions under a certain limit. The appeal and success of this method lies in the fact we circumvent complicated algebro-geometric theory to directly get explicit periodic wave solutions. Moreover, all parameters appearing in the solutions are free variables, whereas usual quasi-periodic solutions involve Riemann constants which are difficult to be determined and need to make complicated Abel transformation on Riemann surface [13-18]. As illustrative examples, we then consider KdV equation and KP equation in this paper. Other many equations also can be dealt with this way.

2. Periodic wave solutions of KdV equation and their reduction

Consider KdV equation

$$u_t + 6uu_x + u_{xxx} = 0. \tag{2.1}$$

Substituting transformation

$$u = u_0 + 2(\ln f)_{xx} \tag{2.2}$$

into Eq.(2.1) and integrating once again, we then get the following bilinear form

$$G(D_x, D_t)f \cdot f = (D_x D_t + 6u_0 D_x^2 + D_x^4 + c)f \cdot f = 0 \tag{2.3}$$

where $c = c(t)$ is a integration constant and u_0 is a constant solution of KdV equation.

The Hirota bilinear differential operator is defined by

$$D_x^m D_t^n f(x,t) \cdot g(x,t) = (\partial_x - \partial_{x'})^m (\partial_t - \partial_{t'})^n f(x,t) g(x,t)|_{x=x',t=t'}$$

The D-operator have good property when acting on exponential functions

$$D_x^m D_t^n e^{\xi_1} \cdot e^{\xi_2} = (k_1 - k_2)^m (\omega_1 - \omega_2)^n e^{\xi_1 + \xi_2}$$

where $\xi_j = k_j x + \omega_j t, j = 1, 2$. More general, we have

$$G(D_x, D_t) e^{\xi_1} \cdot e^{\xi_2} = G(k_1 - k_2, \omega_1 - \omega_2) e^{\xi_1 + \xi_2} \tag{2.4}$$

We consider Riemann theta function solution of KdV equation

$$f = \sum_{n \in Z^N} e^{\pi i < \tau n, n > + 2\pi i < \xi, n >} \tag{2.5}$$

where $n = (n_1, \cdots, n_N), \xi = (\xi_1, \cdots, \xi_N)$, τ is a symmetric matrix and $\mathrm{Im}\tau > 0$, $\xi_j = k_j x + \omega_j t, j = 1, \cdots, N$.

Consider the case when $N = 1$, then (2.5) becomes

$$f = \sum_{n=-\infty}^{\infty} e^{2\pi i n \xi + \pi i n^2 \tau} \tag{2.6}$$

Substituting (2.6) into (2.3) gives

$$\begin{aligned}
Gf\cdot f &= G(D_x,D_t)\sum_{n=-\infty}^{\infty}e^{2\pi in\xi+\pi in^2\tau}\cdot\sum_{m=-\infty}^{\infty}e^{2\pi im\xi+\pi im^2\tau}\\
&=\sum_{n=-\infty}^{\infty}\sum_{m=-\infty}^{\infty}G(D_x,D_t)e^{2\pi in\xi+\pi in^2\tau}\cdot e^{2\pi im\xi+\pi im^2\tau}\\
&=\sum_{n=-\infty}^{\infty}\sum_{m=-\infty}^{\infty}G(2\pi i(n-m)k,2\pi i(n-m)\omega)e^{2\pi i(n+m)\xi+\pi i(n^2+m^2)\tau}\\
&\overset{n+m=m'}{=}\sum_{m'=-\infty}^{\infty}\left\{\sum_{n=-\infty}^{\infty}G(2\pi i(2n-m')k,2\pi i(2n-m')\omega)e^{\pi i[(n^2+(n-m')^2]\tau}\right\}e^{2\pi im'\xi}\\
&\equiv\sum_{m'=-\infty}^{\infty}\bar{G}(m')e^{2\pi im'\xi}=0.
\end{aligned}$$

Noticing that

$$\begin{aligned}
\bar{G}(m') &= \sum_{n=-\infty}^{\infty}G(2\pi i(2n-m')k,2\pi i(2n-m')\omega)e^{\pi i[(n^2+(n-m')^2]\tau}\\
&\overset{n=n'+1}{=}\sum_{n'=-\infty}^{\infty}G(2\pi i[2n'-(m'-2)]k,2\pi i[2n'-(m'-2)]\omega)\\
&\qquad\times\exp(\pi i[(n'^2+(n'-(m'-2))^2]\tau)\exp(2\pi i(m'-1)\tau)\\
&=\bar{G}(m'-2)e^{2\pi i(m'-1)\tau}\\
&=\cdots=\begin{cases}\bar{G}(0)e^{\pi im'(m'-1)}, & m'\ \text{ is }\ \text{even}\\[2ex] \bar{G}(1)e^{\pi i(m'+1)(m'-2)}, & m'\ \text{ is }\ \text{odd}\end{cases}
\end{aligned}$$

which implies that if $\bar{G}(0)=\bar{G}(1)=0$, then

$$\bar{G}(m')=0,\quad m'\in Z$$

In this way, we may let

$$\bar{G}(0)=\sum_{n=-\infty}^{\infty}(-16\pi^2n^2k\omega-96u_0\pi^2k^2n^2+256\pi^4n^4k^4+c)e^{2\pi in^2\tau}=0,\tag{2.7}$$

$$\bar{G}(1)=\sum_{n=-\infty}^{\infty}(-4\pi^2(2n-1)^2k\omega-24u_0\pi^2(2n-1)^2k^2+16\pi^4(2n-1)^4k^4+c)e^{\pi i(2n^2-2n+1)\tau}=0,\tag{2.8}$$

Denote

$$\delta_1(n) = e^{2\pi i n^2 \tau}, \quad \delta_2(n) = e^{\pi i (2n^2-2n+1)\tau},$$

$$a_{11} = -\sum_{n=-\infty}^{\infty} 16\pi^2 n^2 k \delta_1(n), \quad a_{12} = \sum_{n=-\infty}^{\infty} \delta_1(n),$$

$$b_1 = \sum_{n=-\infty}^{\infty} (256\pi^4 n^4 k^4 - 96 u_0 \pi^2 k^2 n^2)\delta_1(n), \quad a_{21} = -\sum_{n=-\infty}^{\infty} 4\pi^2 (2n-1)^2 k \delta_2(n),$$

$$a_{22} = \sum_{n=-\infty}^{\infty} \delta_2(n), \quad b_2 = \sum_{n=-\infty}^{\infty} [16\pi^4 (2n-1)^4 k^4 - 24 u_0 \pi^2 (2n-1)^2 k^2]\delta_2(n),$$

then (2.7) and (2.8) can be written as

$$a_{11}\omega + a_{12}c + b_1 = 0,$$

$$a_{21}\omega + a_{22}c + b_2 = 0.$$

Solving this system, we get

$$\omega = \frac{b_2 a_{12} - b_1 a_{22}}{a_{11}a_{22} - a_{12}a_{21}}, \qquad c = \frac{b_1 a_{21} - b_2 a_{11}}{a_{11}a_{22} - a_{12}a_{21}}. \tag{2.9}$$

Finally we get periodic wave solution

$$u = u_0 + 2(\ln f)_{xx}, \tag{2.10}$$

where f and ω are given by (2.6) and (2.9) respectively. From fig.1-fig.3, we see that the parameter τ does affect the period and shape of wave, but parameter k has effect on the period and shape of wave.

The well-known soliton solution of KdV equation can be obtained as limit of the periodic solution (2.10). For this purpose, we write f as

$$f = 1 + \alpha(e^{2\pi i\xi} + e^{-2\pi i\xi}) + \alpha^4(e^{4\pi i\xi} + e^{-4\pi i\xi}) + \cdots,$$

where $\alpha = e^{i\pi\tau}$.

Setting $u_0 = 0, \quad \xi = \xi' - \tau/2, \quad k' = 2\pi i k, \quad \omega' = 2\pi i\omega$, we get

$$f = 1 + e^{k'x+\omega' t} + \alpha^2 e^{-2\pi i\xi'} + \alpha^2 e^{4\pi i\xi'} + \alpha^6 e^{-4\pi i\xi'} + \cdots$$

$$\longrightarrow 1 + e^{k'x+\omega' t}, \quad \text{as} \quad \alpha \longrightarrow 0$$



Figure 1. One-periodic wave for KdV equation and the effect of parameters on wave shape: (a) along x-axis, (b) along t-axis, where $k = 0.1, \tau = i$.



Figure 2. One-periodic wave for KdV equation and the effect of parameters on wave shape: (a) along x-axis, (b) along t-axis, where $k = 0.1, \tau = 3i$.

So the periodic solution (2.10) convergence to well-known soliton solution

$$u = 2(\ln f)_{xx}, \quad f = 1 + e^{k'x+\omega' t},$$

We only need to prove that

$$\omega' \longrightarrow -k'^3, \quad \text{as} \quad \alpha \longrightarrow 0. \tag{2.11}$$



Figure 3. One-periodic wave for KdV equation and the effect of parameters on wave shape: (a) along x-axis, (b) along t-axis, where $k = 0.15, \tau = i$.

In fact, it is easy to see that

$$\begin{aligned}
a_{11} &= -32k\pi^2(\alpha^2 + 4\alpha^8 + \cdots),\\
a_{12} &= 1 + 2\alpha^2 + \cdots,\\
a_{22} &= 2\alpha + \alpha^5 + \cdots,\\
a_{21} &= -8k\pi^2(\alpha + 9\alpha^5 + \cdots),\\
b_1 &= 256k^4\pi^4(\alpha^2 + 16\alpha^8 + \cdots),\\
b_2 &= 16k^4\pi^4(2\alpha + 81\alpha^5),
\end{aligned}$$

which lead to

$$b_2a_{12} - b_1a_{22} = 32k^4\pi^4\alpha + o(\alpha), \quad a_{11}a_{22} - a_{12}a_{21} = 8k\pi^2\alpha + o(\alpha)$$

Therefore we have

$$\omega \longrightarrow 4k^3\pi^2, \quad \text{as} \ \ \alpha \longrightarrow 0,$$

which implies (2.11).

Now we consider two-periodic wave solution of KdV equation ($N=2$). Substituting (2.5) into (2.3), we have

$$\begin{aligned}
&Gf\cdot f=\sum_{m,n\in Z^2} G(D_x,D_t)e^{2\pi i<\xi,n>+\pi i<\tau n,n>}\cdot e^{2\pi i<\xi,m>+\pi i<\tau m,m>}\\
&=\sum_{m,n\in Z^2} G(2\pi i<n-m,k>,2\pi i<n-m,\omega>)e^{2\pi i<\xi,n+m>+\pi i(<\tau m,m>+<\tau n,n>)}\\
&\overset{n+m=m'}{=}\sum_{m'\in Z^2}\sum_{n_1,n_2=-\infty}^{\infty} G(2\pi i<2n-m',k>,2\pi i<2n-m',\omega>)\\
&\qquad\quad\times\exp(\pi i(<\tau(n-m'),n-m'>+<\tau n,n>))\exp(2\pi i<\xi,m'>)\\
&\equiv\sum_{m'\in Z^2}\bar{G}(m_1',m_2')e^{2\pi i<\xi,m'>}=0.
\end{aligned}$$

It is easy to calculate that

$$\begin{aligned}
&\bar{G}(m_1',m_2')\\
&=\sum_{n_1,n_2=-\infty}^{\infty} G(2\pi i<2n-m',k>,2\pi i<2n-m',\omega>)e^{\pi i(<\tau(n-m'),n-m'>+<\tau n,n>)}\\
&\overset{n_j=n_j'+\delta_{jl},l=1,2}{=}\sum_{n_1,n_2=-\infty}^{\infty} G\left(2\pi i\sum_{j=1}^{2}(2n_j'-(m_j'-2\delta_{jl}))k_j,2\pi i\sum_{j=1}^{2}(2n_j'-(m_j'-2\delta_{jl}))\omega_j\right)\\
&\times\exp\left\{\pi i\sum_{j,k=1}^{2}[(n_j'+\delta_{jl})\tau_{jk}(n_k'+\delta_{kl})+((m_j'-2\delta_{jl}-n_j')+\delta_{jl})\tau_{jk}(m_k'-2\delta_{kl}-n_k')+\delta_{kl})]\right\},\\
&=\begin{cases}\bar{G}(m_1'-2,m_2')e^{2\pi i(m_1'-1)\tau_{11}+2\pi i m_2'\tau_{12}}, & l=1\\[2mm] \bar{G}(m_1',m_2'-2)e^{2\pi i(m_2'-1)\tau_{22}+2\pi i m_1'\tau_{12}}, & l=2\end{cases}
\end{aligned}$$

which implies that if

$$\bar{G}(0,0)=\bar{G}(0,1)=\bar{G}(1,0)=\bar{G}(1,1)=0,$$

then $\bar{G}(m_1',m_2')=0$ and (2.2) is an exact solution of KdV equation.

Denote

$$
\begin{aligned}
&a_{j1} = -\sum_{n_1,n_2=-\infty}^{\infty} 4\pi^2 < 2n - m^j, k > (2n_1 - m_1^j)\delta_j(n),\\
&a_{j2} = -\sum_{n_1,n_2=-\infty}^{\infty} 4\pi^2 < 2n - m^j, k > (2n_2 - m_2^j)\delta_j(n)\\
&a_{j3} = -\sum_{n_1,n_2=-\infty}^{\infty} 24\pi^2 < 2n - m^j, k >^2 \delta_j(n),\\
&a_{j4} = \sum_{n_1,n_2=-\infty}^{\infty} \delta_j(n)\\
&b_j = -\sum_{n_1,n_2=-\infty}^{\infty} 16\pi^4 < 2n - m^j, k >^4 \delta_j(n),\\
&\delta_j(n) = e^{\pi i < \tau(n-m^j), n-m^j > + \pi i < \tau n, n >},\\
&j = 1,2,3,4 \;\; m^1 = (0,0), \quad m^2 = (1,0), \quad m^3 = (0,1), \quad m^4 = (1,1)\\
&A = (a_{kj})_{4\times 4}, \quad b = (b_1, b_2, b_3, b_4)^T,
\end{aligned}
$$

then we have

$$A \begin{pmatrix} \omega_1 \\ \omega_2 \\ u_0 \\ c \end{pmatrix} = b,$$

from which we obtain

$$\omega_1 = \frac{\Delta_1}{\Delta}, \quad \omega_2 = \frac{\Delta_2}{\Delta}, \quad u_0 = \frac{\Delta_3}{\Delta}, \tag{2.12}$$

where $\Delta = |A|$ and $\Delta_1, \Delta_2, \Delta_3$ are produced from Δ by replacing its $1th, 2th, 3th$ column with b respectively. Finally we get two-periodic wave solution (See fig.2)

$$u = u_0 + 2(\ln f)_{xx}, \tag{2.13}$$

where f and ω_1, ω_2 are given by (2.5) and (2.12) respectively. The properties of the solution are shown in fig.4-fig.7. Fig.4 and fig.5 shown effect of parameters k_1, k_2 on period and shape of wave. Fig.7 gives effect of parameter $\tau_{11}, \tau_{12}, \tau_{22}$ on period and shape of wave. The fig.7 and fig.8 show that two-periodic wave can degenerate to one-periodic wave when k_1 is sufficient small.



Figure 4. Two-periodic wave for KdV equation and the effect of parameters on wave shape: (a) along x-axis, (b) along t-axis, where $\tau_{11} = 0.1i, \tau_{12} = 0.2i, \tau_{22} = 3i, k_1 = 0.2, k_2 = -0.3$.



Figure 5. Two-periodic wave for KdV equation and the effect of parameters on wave shape: (a) along x-axis, (b) along t-axis, where $\tau_{11} = 0.1i, \tau_{12} = 0.2i, \tau_{22} = 3i, k_1 = 0.1, k_2 = -0.3$.

The two-soliton solution of KdV equation can be obtained as limit of the periodic solution (2.10). We write f as

$$
\begin{aligned}
&f = 1 + (e^{2\pi i\xi_1} + e^{-2\pi i\xi_1})e^{\pi i\tau_{11}} + (e^{2\pi i\xi_2} + e^{-2\pi i\xi_2})e^{\pi i\tau_{22}} \\
&+(e^{2\pi i(\xi_1+\xi_2)} + e^{-2\pi i(\xi_1+\xi_2)})e^{\pi i(\tau_{11}+2\tau_{12}+\tau_{22})} + \cdots
\end{aligned}
$$

Setting $\xi_1' = 2\pi i\xi_1 - \pi i\tau_{11}, \quad \xi_2' = 2\pi i\xi_2 - \pi i\tau_{22}, \quad \tau_{12} = i\widetilde{\tau}$ ($\widetilde{\tau}$ is a real), we get

$$
\begin{aligned}
&f = 1 + e^{\xi_1'} + e^{\xi_2'} + e^{\xi_1'+\xi_2'+2\pi i\tau_{12}} + \alpha_1^2 e^{-\xi_1'} + \alpha_2^2 e^{-\xi_2'} + \alpha_1^2\alpha_2^2 e^{-\xi_1-\xi_2+2\pi i\tau_{11}} + \cdots \\
&\longrightarrow 1 + e^{\xi_1'} + e^{\xi_2'} + e^{\xi_1'+\xi_2'-2\pi\widetilde{\tau}}, \quad \text{as} \quad \alpha_1, \alpha_2 \longrightarrow 0
\end{aligned}
$$



Figure 6. Two-periodic wave for KdV equation and the effect of parameters on wave shape: (a) along x-axis, (b) along t-axis, where $\tau_{11} = 0.1, \tau_{12} = 0.3i, \tau_{22} = 2i, k_1 = 0.1, k_2 = -0.3$.



Figure 7. Two-periodic wave for KdV equation and the effect of parameters on wave shape: (a) along x-axis, (b) along t-axis, where $\tau_{11} = 0.01, \tau_{12} = 0.2i, \tau_{22} = 3i, k_1 = 0.1, k_2 = -0.2$.

where

$$\alpha_1 = e^{\pi i \tau_{11}}, \quad \alpha_2 = e^{\pi i \tau_{22}}, \quad \xi_i' = k_i' x + \omega_i' t, \quad e^{-2\pi\widetilde{\tau}} = \left(\frac{k_1' - k_2'}{\omega_1' - \omega_2'}\right)^2,$$

$$\omega_i' \to -k_i'^3, \quad \text{as} \quad \alpha_1, \alpha_2 \to 0.$$

3. Periodic wave solutions of KP equation and their reduction

Consider KP equation

$$u_t = (u_{xx} + 3u^2)_x + 3\partial^{-1} u_{yy}. \tag{3.1}$$



Figure 8. Two-periodic wave for KdV equation and the effect of parameters on wave shape: (a) along x-axis, (b) along t-axis, where $\tau_{11} = 0.001, \tau_{12} = 0.2i, \tau_{22} = 3i, k_1 = 0.1, k_2 = -0.2$.

Substituting transformation

$$u = u_0 + 2(\ln f)_{xx} \tag{3.2}$$

into (3.1) and integrating once again, we then get the following bilinear form

$$G(D_x, D_y, D_t) f \cdot f = (D_x D_t - 6u_0 D_x^2 - D_x^4 - 3D_y^2 + c_0) f \cdot f = 0 \tag{3.3}$$

where c_0 is a constant of integration and u_0 is a constant solution of KP equation. Let us consider Riemann theta function solution of KP equation

$$f = \sum_{n \in Z^N} e^{\pi i < \tau n, n > + 2\pi i < \xi, n >} \tag{3.4}$$

where $n = (n_1, \cdots, n_N), \xi = (\xi_1, \cdots, \xi_N)$, τ is a symmetric matrix and $\mathrm{Im}\tau > 0$, $\xi_j = k_j(x + \rho_j y + \omega_j t) + \gamma_j, j = 1, \cdots, N$.

In the following, we consider some special cases. When $N = 1$, (3.4) is reduced to

$$f = \sum_{n=-\infty}^{\infty} e^{2\pi i n \xi + \pi i n^2 \tau} \tag{3.5}$$

Substituting (3.5) into (3.3) gives

$$\begin{aligned}
&Gf\cdot f = G(D_x,D_y,D_t)\sum_{n=-\infty}^{\infty} e^{2\pi in\xi+\pi in^2\tau}\cdot\sum_{m=-\infty}^{\infty} e^{2\pi im\xi+\pi im^2\tau}\\
&=\sum_{n=-\infty}^{\infty}\sum_{m=-\infty}^{\infty} G(D_x,D_y,D_t)e^{2\pi in\xi+\pi in^2\tau}\cdot e^{2\pi im\xi+\pi im^2\tau}\\
&=\sum_{n=-\infty}^{\infty}\sum_{m=-\infty}^{\infty} G(2\pi i(n-m)k,2\pi i(n-m)k\rho,2\pi i(n-m)k\omega)e^{2\pi i(n+m)\xi+\pi i(n^2+m^2)\tau}\\
&\overset{n+m=m'}{=}\sum_{m'=-\infty}^{\infty}\left\{\sum_{n=-\infty}^{\infty} G(2\pi i(2n-m')k,2\pi i(2n-m')k\rho,2\pi i(2n-m')k\omega)e^{\pi i[(n^2+(n-m')^2]\tau}\right\}e^{2\pi im'\xi}\\
&\equiv\sum_{m'=-\infty}^{\infty}\bar{G}(m')e^{2\pi im'\xi}=0.
\end{aligned}$$

Noticing that

$$\begin{aligned}
&\bar{G}(m')=\sum_{n=-\infty}^{\infty} G(2\pi i(2n-m')k,2\pi i(2n-m')k\rho,2\pi i(2n-m')k\omega)e^{\pi i[(n^2+(n-m')^2]\tau}\\
&\overset{n=n'+1}{=}\sum_{n'=-\infty}^{\infty} G(2\pi i[2n'-(m'-2)]k,2\pi i[2n'-(m'-2)]k\rho,2\pi i[2n'-(m'-2)]k\omega)\\
&\qquad\times\exp(\pi i[(n'^2+(n'-(m'-2))^2]\tau)\exp(2\pi i(m'-1)\tau)\\
&=\bar{G}(m'-2)e^{2\pi i(m'-1)\tau}\\
&=\cdots=\begin{cases}\bar{G}(0)e^{\pi im'(m'-1)}, & m' \text{ is even}\\[2ex] \bar{G}(1)e^{\pi i(m'+1)(m'-2)}, & m' \text{ is odd}\end{cases}
\end{aligned}$$

which implies that if $\bar{G}(0)=\bar{G}(1)=0$, then

$$\bar{G}(m')=0,\quad m'\in Z$$

In this way, we may let

$$\bar{G}(0)=\sum_{n=-\infty}^{\infty}(-16\pi^2n^2k^2\omega+96u_0\pi^2k^2n^2-256\pi^4n^4k^4+48\pi^2k^2\rho^2n^2+c_0)e^{2\pi in^2\tau}=0, \tag{3.6}$$

$$\begin{aligned}\bar{G}(1)=\sum_{n=-\infty}^{\infty}&(-4\pi^2(2n-1)^2k^2\omega+24u_0\pi^2(2n-1)^2k^2-16\pi^4(2n-1)^4k^4\\
&+12\pi^2(2n-1)^2k^2\rho^2+c_0)e^{\pi i(2n^2-2n+1)\tau}=0,\end{aligned} \tag{3.7}$$

Denote

$$\alpha = e^{\pi\tau i}, \quad a_{11} = -16\pi^2k^2\sum_{n=-\infty}^{\infty} n^2\alpha^{2n^2}, \quad a_{12} = \sum_{n=-\infty}^{\infty}\alpha^{2n^2},$$

$$b_1 = -\sum_{n=-\infty}^{\infty}(96u_0\pi^2n^2k^2 - 256\pi^4n^4k^4 + 48\pi^2k^2\rho^2n^2)\alpha^{2n^2}, \quad a_{21} = -4\pi^2k^2\sum_{n=-\infty}^{\infty}(2n-1)^2\alpha^{2n^2-2n+1},$$

$$a_{22} = \sum_{n=-\infty}^{\infty}\alpha^{2n^2-2n+1}, \quad b_2 = \sum_{n=-\infty}^{\infty}[16\pi^4(2n-1)^4k^4 - 24u_0\pi^2(2n-1)^2k^2 - 12\pi^2(2n-1)^2k^2\rho^2]\alpha^{2n^2-2n+1},$$

then (3.6) and (3.7) can be written as

$$a_{11}\omega + a_{12}c_0 = b_1,$$

$$a_{21}\omega + a_{22}c_0 = b_2.$$

Solving this system, we get

$$\omega = \frac{b_1a_{22} - b_2a_{12}}{a_{11}a_{22} - a_{12}a_{21}}, \qquad c_0 = \frac{b_2a_{11} - b_1a_{21}}{a_{11}a_{22} - a_{12}a_{21}}. \tag{3.8}$$

Finally we get periodic wave solution of KP equation (3.1)

$$u = u_0 + 2(\ln f)_{xx}, \tag{3.9}$$

where f and ω are given by (3.4) and (3.8) respectively. From fig.9-fig.11, we see that the parameter τ does affect the period and shape of wave, but parameter k has effect on the period and shape of wave.

The well-known soliton solution of the KP equation can be obtained as limit of the periodic solution (3.9). To this end, we write f as

$$f = 1 + \alpha(e^{2\pi i\xi} + e^{-2\pi i\xi}) + \alpha^4(e^{4\pi i\xi} + e^{-4\pi i\xi}) + \cdots.$$

Setting $u_0 = 0,\ \gamma = \gamma_0 - \tau/2,\ k' = 2\pi ik,\ \gamma_0' = 2\pi i\gamma_0$, we get

$$\begin{aligned} f &= 1 + e^{k'(x+\rho y+\omega t)+\gamma_0'} + \alpha^2e^{-k'(x+\rho y+\omega t)-\gamma_0'} + \alpha^2e^{2k'(x+\rho y+\omega t)+2\gamma_0'} \\ &\quad + \alpha^6e^{-2k'(x+\rho y+\omega t)-2\gamma_0'} + \cdots \\ &\longrightarrow 1 + e^{k'(x+\rho y+\omega t)+\gamma_0'}, \quad \text{as} \quad \alpha \longrightarrow 0 \end{aligned}$$

Then the periodic solution (3.9) convergence to well-known soliton solution

$$u = 2(\ln f)_{xx}, \quad f = 1 + e^{k'(x+\rho y+\omega t)+\gamma_0'}. \tag{3.10}$$



Figure 9. One-periodic wave for KP equation and the effect of parameters on wave shape: (a) along x-axis, (b) along y-axis, (b) along t-axis, where $k = 0.1, \tau = i, \rho = 2$.

We only need to prove that

$$\omega \longrightarrow -4k^3\pi^2 + 3k\rho^2, \quad \text{as} \ \ \alpha \longrightarrow 0. \tag{3.11}$$

In fact, it is easy to see that

$$\begin{aligned}
a_{11} &= -32k\pi^2(\alpha^2 + 4\alpha^8 + \cdots), \\
a_{12} &= 1 + 2\alpha^2 + \cdots, \\
a_{21} &= -8\pi^2 k(\alpha + 9\alpha^5 + \cdots), \\
a_{22} &= 2(\alpha + \alpha^5 + \cdots), \\
b_1 &= 512k^4\pi^4(\alpha^2 + 16\alpha^8 + \cdots) - 96k^2\pi^2\rho^2(\alpha^2 + 4\alpha^8 + \cdots), \\
b_2 &= 32k^4\pi^4(\alpha + 81\alpha^5 + \cdots) - 24k^2\pi^2\rho^2(\alpha + 9\alpha^5 + \cdots).
\end{aligned}$$



Figure 10. One-periodic wave for KP equation and the effect of parameters on wave shape: (a) along x-axis, (b) along y-axis, (b) along t-axis, where $k = 0.1, \tau = 3i, \rho = 1$.

Therefore we have

$$b_1 a_{22} - b_2 a_{12} = -32k^4\pi^4\alpha + 24\pi^2 k^2 \rho^2 \alpha + o(\alpha), \quad a_{11}a_{22} - a_{12}a_{21} = 8k\pi^2\alpha + o(\alpha),$$

which lead to (3.11).



Figure 11. One-periodic wave for KP equation and the effect of parameters on wave shape: (a) along x-axis, (b) along y-axis, (c) along t-axis, where $k = 0.2, \tau = i, \rho = 1$.

Now we consider two-periodic wave solution of the KP equation ($N = 2$). Let $\hat{\rho} = (k_1\rho_1, k_2\rho_2)$, $\hat{\omega} = (k_1\omega_1, k_2\omega_2)$. Substituting (3.4) into (3.3), we have

$$
\begin{aligned}
&Gf \cdot f = \sum_{m,n \in Z^2} G(D_x, D_y, D_t) e^{2\pi i <\xi, n> + \pi i <\tau n, n>} \cdot e^{2\pi i <\xi, m> + \pi i <\tau m, m>} \\
&= \sum_{m,n \in Z^2} G(2\pi i < n-m, k >, 2\pi i < n-m, \hat{\rho} >, 2\pi i < n-m, \hat{\omega} >) e^{2\pi i <\xi, n+m> + \pi i (<\tau m, m> + <\tau n, n>)} \\
&\overset{n+m=m'}{=} \sum_{m' \in Z^2} \sum_{n_1, n_2 = -\infty}^{\infty} G(2\pi i < 2n-m', k >, 2\pi i < 2n-m', \hat{\rho} >, 2\pi i < 2n-m', \hat{\omega} >) \\
&\qquad \times \exp(\pi i (< \tau(n-m'), n-m' > + < \tau n, n >)) \exp(2\pi i < \xi, m' >) \\
&\equiv \sum_{m' \in Z^2} \bar{G}(m_1', m_2') e^{2\pi i <\xi, m'>} = 0.
\end{aligned}
$$

It is easy to calculate that

$$\begin{aligned}
&\bar{G}(m_1', m_2') \\
&= \sum_{n_1,n_2=-\infty}^{\infty} G(2\pi i < 2n - m', k >, 2\pi i < 2n - m', \hat{\rho} >, 2\pi i < 2n - m', \hat{\omega} >) e^{\pi i (<\tau(n-m'), n-m'> + <\tau n, n>)} \\
&= \sum_{n_1,n_2=-\infty}^{\infty} G\left(2\pi i \sum_{j=1}^{2} (2n_j' - (m_j' - 2\delta_{jl}))k_j, 2\pi i \sum_{j=1}^{2} (2n_j' - (m_j' - 2\delta_{jl}))k_j \rho_j, 2\pi i \sum_{j=1}^{2} (2n_j' - (m_j' - 2\delta_{jl}))k_j \omega_j \right) \\
&\times \exp\left\{ \pi i \sum_{j,k=1}^{2} [(n_j' + \delta_{jl})\tau_{jk}(n_k' + \delta_{kl}) + ((m_j' - 2\delta_{jl} - n_j') + \delta_{jl})\tau_{jk}(m_k' - 2\delta_{kl} - n_k') + \delta_{kl})] \right\}, \\
&= \begin{cases} \bar{G}(m_1' - 2, m_2') e^{2\pi i (m_1' - 1)\tau_{11} + 2\pi i m_2' \tau_{12}}, & l = 1 \\ \bar{G}(m_1', m_2' - 2) e^{2\pi i (m_2' - 1)\tau_{22} + 2\pi i m_1' \tau_{12}}, & l = 2 \end{cases}
\end{aligned}$$

which implies that if

$$\bar{G}(0,0) = \bar{G}(0,1) = \bar{G}(1,0) = \bar{G}(1,1) = 0,$$

then $\bar{G}(m_1', m_2') = 0$ and (3.2) is an exact solution of the KP equation. Denote

$$\begin{aligned}
&a_{j1} = -4\pi^2 k_1 \sum_{n_1,n_2=-\infty}^{\infty} < 2n - m^j, k > (2n_1 - m_1^j)\delta_j(n), \\
&a_{j2} = -4\pi^2 k_2 \sum_{n_1,n_2=-\infty}^{\infty} < 2n - m^j, k > (2n_2 - m_2^j)\delta_j(n) \\
&a_{j3} = 24\pi^2 \sum_{n_1,n_2=-\infty}^{\infty} < 2n - m^j, k >^2 \delta_j(n), \\
&a_{j4} = \sum_{n_1,n_2=-\infty}^{\infty} \delta_j(n) \\
&b_j = \sum_{n_1,n_2=-\infty}^{\infty} (16\pi^4 < 2n - m^j, k >^4 - 12\pi^2 < 2n - m^j, \hat{\rho} >^2)\delta_j(n), \\
&\delta_j(n) = e^{\pi i <\tau(n-m^j), n-m^j> + \pi i <\tau n, n>}, \quad j = 1, 2, 3, 4 \\
&m^1 = (0,0), \quad m^2 = (1,0), \quad m^3 = (0,1), \quad m^4 = (1,1) \\
&A = (a_{kj})_{4\times 4}, \quad b = (b_1, b_2, b_3, b_4)^T,
\end{aligned}$$

then we have

$$A\begin{pmatrix}\omega_1\\ \omega_2\\ u_0\\ c\end{pmatrix}=b, \tag{3.12}$$

from which we obtain

$$\omega_1=\frac{\Delta_1}{\Delta},\quad \omega_2=\frac{\Delta_2}{\Delta},\quad u_0=\frac{\Delta_3}{\Delta},\quad c_0=\frac{\Delta_4}{\Delta} \tag{3.13}$$

where $\Delta = |A|$ and Δ_j are produced from Δ by replacing its jth column with b respectively. Therefore, we arrive at two-periodic wave solution

$$u=u_0+2(\ln f)_{xx}, \tag{3.14}$$

where f and ω_1, ω_2 are given by (3.4) and (3.13) respectively. The properties of the solution are shown in fig.12-fig.16. Fig.12 and fig.13 shown effect of parameters k_1, k_2 on period and shape of wave. Fig.14 gives effect of parameter $\tau_{11}, \tau_{12}, \tau_{22}$ on period and shape of wave. The fig.15 and fig.16 show that two-periodic wave can degenerate to one-periodic wave when k_1 is sufficient small.

Two-soliton solution of the KP equation can be obtained as limit of the periodic solution (3.14). We write f as

$$\begin{aligned}f&=1+(e^{2\pi i\xi_1}+e^{-2\pi i\xi_1})e^{\pi i\tau_{11}}+(e^{2\pi i\xi_2}+e^{-2\pi i\xi_2})e^{\pi i\tau_{22}}\\&+(e^{2\pi i(\xi_1+\xi_2)}+e^{-2\pi i(\xi_1+\xi_2)})e^{\pi i(\tau_{11}+2\tau_{12}+\tau_{22})}+\cdots\end{aligned}$$

Setting $\gamma_1'=\gamma_1+\frac{1}{2}\tau_{11},\ \gamma_2'=\gamma_2+\frac{1}{2}\tau_{22},\ \xi_1'=2\pi i\xi_1-\pi i\tau_{11},\ \ \xi_2'=2\pi i\xi_2-\pi i\tau_{22},\ \ \tau_{12}=i\tau'$ (τ' is a real), we get

$$\begin{aligned}f&=1+e^{\xi_1'}+e^{\xi_2'}+e^{\xi_1'+\xi_2'+2\pi i\tau_{12}}+\alpha_1^2e^{-\xi_1'}+\alpha_2^2e^{-\xi_2'}+\alpha_1^2\alpha_2^2e^{-\xi_1-\xi_2+2\pi i\tau_{11}}+\cdots\\&\longrightarrow 1+e^{\xi_1'}+e^{\xi_2'}+e^{\xi_1'+\xi_2'-2\pi\tau'},\quad \text{as}\ \ \alpha_1,\alpha_2\longrightarrow 0\end{aligned}$$

where

$$\alpha_1=e^{\pi i\tau_{11}},\quad \alpha_2=e^{\pi i\tau_{22}},\quad \xi_j'=2\pi i[k_j(x+\rho_j+\omega_jt)+\gamma_j'],$$

$$e^{-2\pi\tau'}=\frac{(k_1-k_2)^2-(\rho_1-\rho_2)^2}{(k_1+k_2)^2-(\rho_1-\rho_2)^2},\quad \omega_i'\to-4k_j^3\pi^2+3k\rho_j^2,\qquad \text{as}\qquad \alpha_1,\alpha_2\to 0.$$



Figure 12. Two-periodic wave for KP equation and the effect of parameters on wave shape: (a) along x-axis, (b) along y-axis, (c) along t-axis, where $k_1 = 0.2, k_2 = -0.3$, $\tau_{11} = 0.1i, \tau_{12} = 0.2i, \tau_{22} = 3i, \rho_1 = 1, \rho_2 = 2$.

Acknowledgments

The work described in this paper was partially supported by a grant from City University of Hong Kong (Project No. 9040967).



Figure 13. Two-periodic wave for KP equation and the effect of parameters on wave shape: (a) along x-axis, (b) along y-axis, (c) along t-axis, where $k_1 = 0.1, k_2 = -0.3$, $\tau_{11} = 0.1i, \tau_{12} = 0.2i, \tau_{22} = 3i, \rho_1 = 1, \rho_2 = 2$.

References

[1] M.J. Ablowitz and P.A. Clarkson, Solitons, Nonlinear Evolution Equations and Inverse Scattering, Cambridge University Press, 1991.

[2] R. Beals and R.R. Coifman, Comm. pure. Appl. Math. 37 (1984) 39.

[3] V.B. Matveev and M.A. Salle, Darboux transformation and solitons, Springer, Berlin, 1991.

[4] C.H. Gu, H.S. Hu and Z.X. Zhou, Darboux Transformations in Soliton Theory and its Geometric Applications,Shanghai Science Technology Publisher, Shanghai, 1999.

[5] S.B. Leble and N.V. Ustinov, J. Phys. A 26 (1993) 5007.

[6] P.G. Esteevez, J. Math. Phys. 40 (1999) 1406.

[7] R. Hirota and J. Satsuma, Prog. Theor. Phys. **57** (1977) 797.

[8] R. Hirota, Direct methods in soliton theory, Springer, 2004.



Figure 14. Two-periodic wave for KP equation and the effect of parameters on wave shape: (a) along x-axis, (b) along y-axis, (c) along t-axis, where $k_1 = 0.2, k_2 = -0.3$, $\tau_{11} = 0.1i, \tau_{12} = 0.2i, \tau_{22} = i, \rho_1 = 0.2, \rho_2 = 0.3$.

[9] X. B. Hu, P. A. Clarkson, J. Phys. A,**28**(1995), 5009.

[10] X. B. Hu, C X Li, J.J. C. Nimmo, G. F. Yu, J. Phys A, **38**(2005), 195.

[11] R Hirota, Y. Ohta, J. Phys Soc. Jpn **60**(1991) 798

[12] J X Zhao, C X Li, X. B Hu, J. Phys Soc. Jpn **73**(2004) 1159

[13] B. A. Dubrovin, Funct. Anal. Appl. **9** (1975) 41.

[14] A. Its and V. Matveev, Funct. Anal. Appl. **9** (1975) 69.

[15] E. Belokolos, A. Bobenko, V. Enol'skij, A, Its and V. Matveev, *Algebro-Geometrical Approach to Nonlinear Integrable Equations*, Springer, Berlin, 1994.

[16] S.P. Novikov, and S.V. Manakov, L.P. Pitaeskii and V.E. Zakharov, *Theory of Solitons, The Snverse Scattering Methods*, Nauka, Moscow, 1980.

[17] P. L. Christiansen, J. C. Eilbeck, V. Z. Enolskii and N. A. Kostov, Proc. R. Soc. London A Math. **451** (1995) 685



Figure 15. Two-periodic wave for KP equation and the effect of parameters on wave shape: (a) along x-axis, (b) along y-axis, (c) along t-axis, where $k_1 = 0.01, k_2 = -0.3$, $\tau_{11} = 0.1i, \tau_{12} = 0.2i, \tau_{22} = i, \rho_1 = 0.2, \rho_2 = 0.3$.

[18] R. G. Zhou, J. Math. Phys. **38** (1997) 2535.

[19] C. W. Cao, Y. T. Wu and X. G. Geng, J. Math. Phys. **40** 1999, 3948.

[20] X. G. Geng and Y. T. Wu, J. Math. Phys. **40** (1999), 2971.

[21] X. G. Geng, C. W. Cao and H. H. Dai, J. Phys. A, **34** 2001, 989.

[22] Z. J. Qiao, Reviews in Math. Phys. **13** (2001), 545.

[23] C. W. Cao, X. G. Geng and H. Y. Wang, J. Math. Phys. **43** 2002, 621.

[24] H. H. Dai and X. G. Geng, Chaos, Solitons and Fractrals, **18**(2003), 1031

[25] B. Deconinck, M. V. Hoeij, Physica D, **152-153**(2001) 28

[26] B Docninck, M Meil, A. Bobenko, M. V. Hoeij and M. Schmies, Math. of Comput. 1417.



Figure 16. Two-periodic wave for KP equation and the effect of parameters on wave shape: (a) along x-axis, (b) along y-axis, (c) along t-axis, where $k_1 = 0.001, k_2 = -0.3$, $\tau_{11} = 0.1i, \tau_{12} = 0.2i, \tau_{22} = i, \rho_1 = 0.2, \rho_2 = 0.3$.